

Character sums of composite moduli and hybrid subconvexity

Roman Holowinsky, Ritabrata Munshi, and Zhi Qi

On the occasion of James W. Cogdell's 60th birthday.

ABSTRACT. Let $M = M_1 M_2 M_3$ be the product of three distinct primes and let $\chi = \chi_1 \chi_2 \chi_3$ be a Dirichlet character of modulus M such that each χ_i is a primitive character modulo M_i for $i = 1, 2, 3$. In this paper, we provide a δ -symbol method for obtaining non-trivial cancellation in smooth character sums of the form $\sum_{n=1}^{\infty} \chi(n) W(n/N)$, with N roughly of size $\sqrt{M}$ and W a smooth compactly supported weight function on $(0, \infty)$. As a corollary, we establish hybrid subconvexity bounds for the associated Dirichlet L -function.

1. Introduction and main results

Let $L(s, \pi)$ be the L -function associated with an irreducible cuspidal automorphic representation π with unitary central character. Analysis of $L(s, \pi)$ leads to information about the arithmetic or algebraic structure associated with π and non-trivial estimates for $L(s, \pi)$ in terms of its analytic conductor $Q(s, \pi)$ (for values of s in the critical strip $0 \leq \Re(s) \leq 1$) often result in non-trivial applications. One classical problem, the subconvexity problem, is to establish a bound of the form

$$L(s, \pi) \ll Q(s, \pi)^{1/4-\delta}$$

for some $\delta > 0$ when $\Re(s) = 1/2$. In general, for $\Re(s) = 1/2$, one has the convexity bound

$$L(s, \pi) \ll_{\varepsilon} Q(s, \pi)^{1/4+\varepsilon}$$

while the Riemann Hypothesis for $L(s, \pi)$ would imply the Lindelöf Hypothesis

$$L(s, \pi) \ll_{\varepsilon} Q(s, \pi)^{\varepsilon}.$$

Although the convexity bound is far from the expected Lindelöf bound, any power saving in the conductor is often sufficient for applications. For example, subconvexity for $L(\frac{1}{2} + it, \text{Sym}^2 f)$ and $L(\frac{1}{2}, \text{Sym}^2 f \times \varphi)$, where t is a fixed real number, f is a varying holomorphic eigencuspform (with $\text{Sym}^2 f$ its symmetric square) and

1991 *Mathematics Subject Classification.* 11M06.

Key words and phrases. Dirichlet L -functions, subconvexity, character sums.

The first author completed this work through the support of the NSF grant DMS-1068043. The second author was partly supported by Swarna Jayanti Fellowship, 2011-12, DST, Govt. of India.

φ is a fixed Hecke-Maass eigencuspform for the modular group $\mathrm{SL}_2(\mathbb{Z})$, implies the Mass Equidistribution Conjecture (a holomorphic analogue of the Quantum Unique Ergodicity Conjecture [RS94]) for $\mathrm{SL}_2(\mathbb{Z})$. The subconvexity problem has thus received much attention in various settings recently, however, a general method of proof for all π does not yet exist.

In a collection of works by the authors, see for example [Mun14a], [Mun14b], [Mun13a], [Mun13b], [HM12], and [HMQ14], several methods have been developed to investigate the subconvexity problem particularly in the case of Rankin-Selberg convolution L -functions where multiple parameters are varying. Such methods have led to a variety of hybrid subconvexity results, most recently demonstrating that subconvexity bounds are more readily obtained for $L(\frac{1}{2}, \mathrm{Sym}^2 f \times \varphi)$ when both f and φ are varying. Indeed, in [HMQ14] the authors establish the following result.

THEOREM. *Suppose $k > \kappa \geq 2$ are integers, with k even, P is a prime, f is a Hecke cusp form of weight k for $\mathrm{SL}(2, \mathbb{Z})$, and g a newform of weight 2κ and level P . Then we have*

$$L\left(\frac{1}{2}, \mathrm{Sym}^2 f \otimes g\right) \ll_{\varepsilon, \kappa} \begin{cases} k^{13/29} P^{25/29} (kP)^\varepsilon, & \text{if } P^{13/64} < k \leq P^{4/13}, \\ (P + k^{13/7} P^{3/7}) (kP)^\varepsilon, & \text{if } P^{4/13} < k < P^{3/8}. \end{cases}$$

This bound beats the convexity bound $kP^{3/4} (kP)^\varepsilon$ when $P^{13/64+\delta} < k < P^{3/8-\delta}$ for some $0 < \delta < 11/128$.

Although more parameters are contributing to the complexity and analytic conductor of the L -function in such hybrid subconvexity problems, these situations are amenable to a larger collection of analytic tools and methods. For example, if $\pi = f_1 \times f_2$ with each f_i a holomorphic newform of varying level N_i and $(N_1, N_2) = 1$, then one has several natural “families” and “sub-families” of L -functions to which $L(s, \pi)$ might be associated. In order to prove subconvexity for $L(s, f_1 \times f_2)$, one might choose to first study a moment average over a basis of newforms of level N_1 , of level N_2 , or average over both N_1 and N_2 . If, instead, only one of the levels is varying, then we immediately lose that additional degree of freedom.

Such hybrid subconvexity problems therefore raise a question regarding structure and which family/moment of L -functions one should consider. In order to establish subconvexity in the case of $\pi = f_1 \times f_2$ above, it was seen in [HM12] that one should average over the larger level family when studying a second moment while one should average over the smaller level family, as in [HT14], when studying a first moment. If one were to study the first moment over the larger level family, then one obtains exact evaluations of the moment average rather than subconvexity (see for example [MR12], [FW09], [Nel13]). Ultimately, the subconvexity problem boils down to having a sufficient number of points of summation relative to the conductor and complexity of the L -function one is considering, without having too many points of summation.

In an attempt to better understand the underlying structure of such hybrid subconvexity results, we turn to the classical example of $\mathrm{GL}(1)$ convolutions. Of course, one has Burgess’ well known result for Dirichlet L -functions of a primitive character χ of modulus M ([Bur63, Theorem 3]),

$$L\left(\frac{1}{2} + it, \chi\right) \ll_{\varepsilon, t} M^{3/16+\varepsilon}.$$

But this does not close the subject. For example, recent work by Milićević [Mil14] on powerful moduli, improves on Burgess' bound for Dirichlet characters with moduli a sufficiently large power of a prime.

In this paper, we present a method for obtaining subconvexity results when the modulus of the Dirichlet character is “moderately” composite. Specifically, when the modulus is a product of three distinct primes $M = M_1 M_2 M_3$. Our method is an adaptation of the one presented in [Mun14b]. Since we are dealing only with Dirichlet characters, the method becomes more transparent. However, our main result is weaker compared to the Burgess bound. As such, this paper does not prove any new result and one should view this work mainly as pedagogical. Our method easily generalizes in the case of “highly” composite moduli. With the availability of more factors, one has more options to design a “conductor lowering” mechanism. Nevertheless, we feel that in this case, the q -analogue of the van der Corput method (see Theorem 12.13 of [IK04]) is much stronger.

Recall that Burgess' bound for Dirichlet L -functions relies on the estimation of the character sum ([Bur62a, Theorem 1], [Bur63, Theorem 2]),

$$\sum_{n=N+1}^{N+H} \chi(n) \ll_{\varepsilon} H^{1/2} N^{3/16+\varepsilon}.$$

When χ is of prime modulus p , the proof of this bound in [Bur62b, Bur63] features the application of an important estimate of Weil for $\sum_{x \in \mathbb{F}_p^*} \chi(f(x))$.

For our purpose we shall consider the smooth character sum

$$(1.1) \quad S_{\chi}(N) := \sum_{n=1}^{\infty} \chi(n) W\left(\frac{n}{N}\right),$$

where W is a smooth weight function on $(0, \infty)$ supported in the interval $[1, 2]$ and satisfying $W^{(j)}(x) \ll_j 1$. We shall obtain the following result on this smooth character sum. Interestingly, our proof also depends on a certain bound due to Deligne and Fu for $\sum_{(x_1, x_2) \in \mathbb{F}_p^{*2}} \chi(x_1 x_2) e(f(x_1, x_2)/p)$ which is rooted in algebraic geometry over a finite field like Weil's bound.

THEOREM 1. *Let M_1, M_2, M_3 be three distinct primes and set $M := M_1 M_2 M_3$. Let χ_i be a primitive character modulo M_i and set $\chi := \chi_1 \chi_2 \chi_3$. For*

$$M_1 \ll N \ll M_1 \min \{M_2^{2/3}, M_3^2\}$$

we have

$$(1.2) \quad S_{\chi}(N) \ll_{\varepsilon} \left(M_2^{1/2} M_3^{1/2} + M_1^{1/4} M_2^{1/2} N^{1/4} + M_3^{1/2} N^{3/4} \right) M^{\varepsilon}.$$

As a corollary, we get the following hybrid subconvexity result.

COROLLARY 1. *Let M_i, M, χ_i and χ , for $i = 1, 2, 3$, be given as in Theorem 1. Set $\theta_i := \log M_i / \log M$ so that $\theta_1 + \theta_2 + \theta_3 = 1$. If*

$$(1.3) \quad 2\delta \leq \theta_1 \leq \frac{1}{2} - \delta, \quad \theta_3 \leq \frac{1}{4} - \delta, \quad \frac{1}{2} + 3\delta \leq \theta_1 + 2\theta_3$$

for some $\delta > 0$, then

$$L\left(\frac{1}{2}, \chi\right) \ll_{\varepsilon} M^{\frac{1}{4} - \frac{\delta}{2} + \varepsilon}.$$

EXAMPLE. *When $(\theta_1, \theta_2, \theta_3) = (\frac{5}{12}, \frac{5}{12}, \frac{1}{6})$, one may choose $\delta = \frac{1}{12}$ so that in this case*

$$L\left(\frac{1}{2}, \chi\right) \ll M^{\frac{5}{24} + \varepsilon}.$$

REMARK. *Of course, the set of triples $(\theta_1, \theta_2, \theta_3)$ for which one obtains non-trivial estimates for $S_\chi(N)$ (and therefore subconvexity bounds for the corresponding set of $L(\frac{1}{2}, \chi)$) can be extended upon permuting the subscripts 1, 2, 3.*

We shall see that a moment average will not be necessary in establishing Theorem 1. Instead, the appropriate number of points of summation will be introduced directly via a δ -symbol method which we describe in the next section. A similar method may be found in [Mun14b]. Furthermore, such a technique with similar arguments would establish subconvexity in the case of $\chi = \chi_1 \chi_2$ when $\chi_1 = M_1$ and $\chi_2 = M_2^2$ and seemingly in higher rank cases when the conductor is of an appropriate form. However, we do not yet see a blanket general structure to classify all situations in which such a δ -symbol method would establish subconvexity.

2. Preliminaries

2.1. Dirichlet L -functions and character sums. For a positive integer $M \geq 2$ let χ be a primitive Dirichlet character of modulus M . The Dirichlet L -function for χ is given by

$$L(s, \chi) = \sum_{n=1}^{\infty} \frac{\chi(n)}{n^s},$$

where the series converges for $\Re(s) > 0$.

From the approximate functional equation and a dyadic partition of unity, one has

$$L\left(\frac{1}{2}, \chi\right) \ll_{\varepsilon} M^{\varepsilon} \sum_N \frac{|S_\chi(N)|}{\sqrt{N}} \left(1 + \frac{N}{\sqrt{M}}\right)^{-A},$$

where $A > 0$, N ranges over $2^{\nu/2}$ for $\nu = -1, 0, 1, 2, \dots$, and $S_\chi(N)$ is the smooth character sum associated to χ defined in (1.1) for some weight function W .

The contribution from those $N \gg M^{1/2+\varepsilon}$ is made negligible by choosing A above to be sufficiently large. Trivially, $|S_\chi(N)| \ll N$, and therefore if $N \ll M^{1/2-\delta}$ for a given $\delta > 0$ then $|S_\chi(N)|/\sqrt{N} \ll M^{1/4-\delta/2}$. Hence we are left with

$$(2.1) \quad L\left(\frac{1}{2}, \chi\right) \ll_{\varepsilon} \left(M^{1/4-\delta/2} + \max_{M^{1/2-\delta} \ll N \ll M^{1/2+\varepsilon}} \frac{|S_\chi(N)|}{\sqrt{N}} \right) M^{\varepsilon}.$$

Thus subconvexity bounds will now follow if one is able to non-trivially bound $S_\chi(N)$ for $M^{1/2-\delta} \ll N \ll M^{1/2+\varepsilon}$.

2.2. A modified δ -symbol method. One of our main analytic tools for the proof of Theorem 1 will be a version of the circle method introduced in [DFI93] and [HB96]. We start with a smooth approximation of the δ -symbol as described in [HB96].

LEMMA 2. *For any $Q > 1$ there is a positive constant c_Q , and a smooth function $h(x, y)$ defined on $(0, \infty) \times (-\infty, \infty)$, such that*

$$(2.2) \quad \delta(n, 0) = \frac{c_Q}{Q^2} \sum_{q=1}^{\infty} \sum_{a \pmod{q}}^* e\left(\frac{an}{q}\right) h\left(\frac{q}{Q}, \frac{n}{Q^2}\right).$$

The constant c_Q satisfies $c_Q = 1 + O_A(Q^{-A})$ for any $A > 0$. Moreover $h(x, y) \ll 1/x$ for all y , and $h(x, y)$ is non-zero only for $x \leq \max\{1, 2|y|\}$.

The smooth function $h(x, y)$ satisfies (see [HB96])

$$(2.3) \quad x^i \frac{\partial^i h}{\partial x^i}(x, y) \ll_i \frac{1}{x} \quad \text{and} \quad \frac{\partial h}{\partial y}(x, y) = 0$$

for $x \leq 1$ and $|y| \leq x/2$. Also for $|y| > x/2$, we have

$$(2.4) \quad x^i y^j \frac{\partial^{i+j} h}{\partial x^i \partial y^j}(x, y) \ll_{i,j} \frac{1}{x}.$$

Our variant of the δ -method makes use of the following observation,

$$\delta(n, 0) = \mathbb{1}_{K|n} \delta(n/K, 0),$$

where K is a positive integer and $\mathbb{1}_{K|n}$ is equal to 1 or 0 according as $K|n$ or not. From this (2.2) may be written in the following form upon detecting the condition $K|n$ with additive characters.

$$(2.5) \quad \delta(n, 0) = \frac{c_Q}{KQ^2} \sum_{q=1}^{\infty} \sum_{a \pmod{q}}^* \sum_{b \pmod{K}} e\left(\frac{an}{qK}\right) e\left(\frac{bn}{K}\right) h\left(\frac{q}{Q}, \frac{n}{KQ^2}\right).$$

2.3. Deligne Bound for Character Sums. Suppose p is a prime and χ is a primitive character modulo p . Let $m, n \in \mathbb{F}_p$. Consider the sum

$$(2.6) \quad \mathfrak{S}_{\chi}(m, n) = \sum_{x_1 \in \mathbb{F}_p^*} \chi(x_1) \overline{\chi}(m + x_1) e\left(\frac{nx_1}{p}\right).$$

Clearly, we have $\mathfrak{S}_{\chi}(0, 0) = p - 1$. In the case $m = 0$ and $n \neq 0$ the character sum reduces to

$$\mathfrak{S}_{\chi}(0, n) = \sum_{x_1 \in \mathbb{F}_p^*} e\left(\frac{nx_1}{p}\right) = -1.$$

In the case $n = 0$ and $m \neq 0$ we have

$$\mathfrak{S}_{\chi}(m, 0) = \sum_{x_1 \in \mathbb{F}_p^*} \overline{\chi}(m\overline{x}_1 + 1) = -1.$$

Finally, we suppose $mn \neq 0$. In this case we will use the following relation

$$\overline{\chi}(a) g_{\chi} = \sum_{b \in \mathbb{F}_p^*} \chi(b) e\left(\frac{ab}{p}\right),$$

which holds for any $a \in \mathbb{F}_p$. Here

$$g_{\chi} = \sum_{a \in \mathbb{F}_p^*} \chi(a) e\left(\frac{a}{p}\right)$$

is the Gauss sum associated with the character χ . Using this relation we rewrite the above character sum as

$$(2.7) \quad \mathfrak{S}_{\chi}(m, n) = \frac{1}{g_{\chi}} \sum_{\mathbf{x} \in \mathbb{F}_p^{*2}} \chi(x_1) \chi(x_2) e\left(\frac{f(\mathbf{x})}{p}\right),$$

where

$$f(\mathbf{x}) = nx_1 + mx_2 + x_1x_2.$$

In order to obtain a non-trivial estimate, we analyse the sum (2.7) using Deligne's work as has been developed in [Fu09]. Let us briefly recall the main result of [Fu09] concerning sums of the form

$$\sum_{\mathbf{x} \in \mathbb{F}_p^{*r}} \chi_1(x_1) \chi_2(x_2) \dots \chi_r(x_r) \psi(f(\mathbf{x})),$$

where ψ is a non-trivial additive character modulo p . Here

$$f(\mathbf{x}) = \sum_{\mathbf{i} \in \mathbb{Z}^r} a_{\mathbf{i}} \mathbf{x}^{\mathbf{i}}$$

is a Laurent polynomial with coefficients $a_{\mathbf{i}} \in \mathbb{F}_p$. Let $\Delta_{\infty}(f)$ be the Newton polyhedron associated with f . This is given by the convex hull in $\mathbb{R}^r$ of the set

$$\{\mathbf{i} \in \mathbb{Z}^r : a_{\mathbf{i}} \neq 0\} \cup \{\mathbf{0}\}.$$

The Laurent polynomial f is said to be non-degenerate with respect to $\Delta_{\infty}(f)$ if for any face τ of $\Delta_{\infty}(f)$ not containing the origin, the locus

$$\frac{\partial f_{\tau}}{\partial x_1} = \dots = \frac{\partial f_{\tau}}{\partial x_r} = 0$$

in the torus $\mathbb{T}_{\mathbb{F}_p}^r = \mathbb{F}_p^{*r}$ is empty, where f_{τ} denotes the sub-polynomial

$$f_{\tau}(\mathbf{x}) = \sum_{\mathbf{i} \in \tau} a_{\mathbf{i}} \mathbf{x}^{\mathbf{i}}.$$

If $\dim \Delta_{\infty}(f) = r$ and f is non-degenerate with respect to $\Delta_{\infty}(f)$, then we have

$$(2.8) \quad \sum_{\mathbf{x} \in \mathbb{F}_p^{*r}} \chi_1(x_1) \chi_2(x_2) \dots \chi_r(x_r) \psi(f(\mathbf{x})) \ll p^{r/2},$$

where the implied constant is independent of p .

Let us now return to the special case of (2.7) with $nm \neq 0$. The Newton polyhedron $\Delta_{\infty}(f)$ of f is given by the convex hull of

$$\{\mathbf{0}, \mathbf{e}_1, \mathbf{e}_2, \mathbf{e}_1 + \mathbf{e}_2\},$$

which is 2 dimensional. Here $\mathbf{e}_1 = (1, 0)$ and $\mathbf{e}_2 = (0, 1)$ are the standard basis vectors. We have

$$\frac{\partial f}{\partial x_1} = n + x_2, \quad \frac{\partial f}{\partial x_2} = m + x_1.$$

Let g be a sub-polynomial of f such that the equations $\partial g / \partial x_1(\mathbf{x}) = \partial g / \partial x_2(\mathbf{x}) = 0$ are solvable on $\mathbb{F}_p^{*2}$. It is easy to verify that one must have $g = 0$ or $g = f$. It is clear that neither 0 nor f is equal to f_{τ} for any face τ of $\Delta_{\infty}(f)$ not containing the origin. This proves that f is non-degenerate with respect to $\Delta_{\infty}(f)$. Using (2.8) along with the expression (2.7) of $\mathfrak{S}_{\chi}(m, n)$, we obtain

$$\mathfrak{S}_{\chi}(m, n) \ll \sqrt{p}.$$

We have arrived at the following Lemma.

LEMMA 3. *Let p be a prime and χ be a primitive character modulo p . For $m, n \in \mathbb{F}_p$ define the character sum $\mathfrak{S}_{\chi}(m, n)$ by (2.6). Then we have*

- $\mathfrak{S}_{\chi}(0, 0) = p - 1$,
- $\mathfrak{S}_{\chi}(m, n) = -1$ if $nm = 0$ and either $m \neq 0$ or $n \neq 0$, and
- $\mathfrak{S}_{\chi}(m, n) \ll \sqrt{p}$ if $mn \neq 0$.

3. Proof of Theorem 1 and Corollary 1

Let M_i , with $i = 1, 2, 3$, be three distinct primes, and set $M = M_1 M_2 M_3$. Let χ_i be primitive characters modulo M_i and set $\chi := \chi_1 \chi_2 \chi_3$. Suppose W is a real-valued smooth function on $(0, \infty)$ supported in $[1, 2]$ and satisfying

$$W^{(j)}(x) \ll_j 1.$$

We shall consider the smooth character sum

$$S_\chi(N) = \sum_{n \in \mathbb{Z}} \chi_1 \chi_2 \chi_3(n) W\left(\frac{n}{N}\right)$$

when $M_1 \ll N \ll M_1 \min\{M_2^{2/3}, M_3^2\}$ (conditions which arise in the course of the proof). Our goal is to establish a non-trivial bound which will be used in application to the subconvexity problem.

3.1. Applying the δ -method. We first write

$$S_\chi(N) = \sum_{n, m \in \mathbb{Z}} \chi_1 \chi_2(n) \chi_3(m) \delta(n - m, 0) W\left(\frac{n}{N}\right) V\left(\frac{m}{N}\right),$$

where V is a smooth function with support $[1/2, 3]$ and such that $V(x) = 1$ for $x \in [1, 2]$ with $V^{(j)}(x) \ll_j 1$.

We apply the modified δ -symbol method described in Section 2.2, with the divisibility modulus $K = M_1$ and $Q = \sqrt{N/M_1}$. From (2.5) we get

$$S_\chi(N) = \frac{c_Q}{N} \sum_{n, m \in \mathbb{Z}} \chi_1 \chi_2(n) \chi_3(m) \sum_{q=1}^{\infty} \sum_{a \pmod{q}}^* \sum_{b \pmod{M_1}}^* e\left(\frac{a(n-m)}{qM_1}\right) e\left(\frac{b(n-m)}{M_1}\right) h\left(\frac{q}{Q}, \frac{n-m}{N}\right) W\left(\frac{n}{N}\right) V\left(\frac{m}{N}\right).$$

In order to have enough points of summation, it is required that

$$(3.1) \quad M_1 \ll N.$$

Estimating trivially at this stage we get $S_\chi(N) \ll N^2$. So our job is to save more than N .

3.2. Poisson summation.

3.2.1. *Poisson summation in the m -sum.* Poisson summation over m gives

$$\begin{aligned} \sum_{m \in \mathbb{Z}} \chi_3(m) e\left(-\frac{am}{qM_1} - \frac{bm}{M_1}\right) h\left(\frac{q}{Q}, \frac{n-m}{N}\right) V\left(\frac{m}{N}\right) \\ = \frac{N}{qM_1 M_3} \sum_{m \in \mathbb{Z}} \mathfrak{C}(m, q, a, b) \mathcal{J}(m, n, q) \end{aligned}$$

where the character sum is given by

$$\mathfrak{C}(m, q, a, b) = \sum_{c \pmod{qM_1 M_3}} \chi_3(c) e\left(-\frac{ac}{qM_1} - \frac{bc}{M_1} + \frac{mc}{qM_1 M_3}\right),$$

and the integral is

$$\mathcal{J}(x, v, q) = \int_{\mathbb{R}} h\left(\frac{q}{Q}, \frac{v}{N} - u\right) V(u) e\left(-\frac{Nxu}{qM_1 M_3}\right) du.$$

Applying integration-by-parts and the bounds from (2.4), we see that if $|m| \gg QM_1M_3M^\varepsilon/N$ then the integral is negligibly small (i.e. $O_A(M^{-A})$ for any $A > 0$).

We impose the restriction that $Q \ll M_3$ with a sufficiently small implied constant. This is equivalent to having

$$(3.2) \quad N \ll M_1M_3^2$$

Under this condition we have $(q, M_3) = 1$, and hence the character sum $\mathfrak{C}(m, q, a, b)$ splits as

$$\sum_{c \pmod{qM_1}} e\left(-\frac{ac}{qM_1} - \frac{bc}{M_1} + \frac{\overline{M_3}mc}{qM_1}\right) \sum_{d \pmod{M_3}} \chi_3(d) e\left(\frac{\overline{qM_1}md}{M_3}\right).$$

It vanishes save for $(a + qb)M_3 \equiv m \pmod{qM_1}$ and $(m, M_3) = 1$ in which case we have

$$qM_1\varepsilon_3\sqrt{M_3}\chi_3(qM_1)\overline{\chi}_3(m).$$

(Here ε_i stands for the sign of the Gauss sum associated with the character χ_i , i.e. $g_{\chi_i} = \varepsilon_i\sqrt{M_i}$.) Observe that the congruence condition above implies $(m, q) = 1$. It follows that

$$\begin{aligned} S_\chi(N) &= \frac{\varepsilon_3 c Q}{\sqrt{M_3}} \sum_{q \ll Q} \sum_{\substack{|m| \ll QM_1M_3M^\varepsilon/N \\ (m, qM_3)=1}} \chi_3(qM_1)\overline{\chi}_3(m) \sum_{n \in \mathbb{Z}} \chi_1\chi_2(n) \\ &\quad \sum_{\substack{a \pmod{q} \\ (a+qb)M_3 \equiv m \pmod{qM_1}}}^* \sum_{b \pmod{M_1}} e\left(\frac{an}{qM_1} + \frac{bn}{M_1}\right) \mathcal{J}(m, n, q) W\left(\frac{n}{N}\right) + O_A(M^{-A}). \end{aligned}$$

We thus need to consider the sum

$$\begin{aligned} &\frac{1}{\sqrt{M_3}} \sum_{q \ll Q} \sum_{\substack{|m| \ll QM_1M_3M^\varepsilon/N \\ (m, qM_3)=1}} \chi_3(q)\overline{\chi}_3(m) \\ &\quad \sum_{n \in \mathbb{Z}} \chi_1\chi_2(n) e\left(\frac{\overline{M_3}mn}{qM_1}\right) \mathcal{J}(m, n, q) W\left(\frac{n}{N}\right). \end{aligned}$$

At this stage trivial estimation gives

$$S_\chi(N) \ll N\sqrt{M_3}M^\varepsilon,$$

and it remains to save more than $\sqrt{M_3}$.

3.2.2. Poisson summation in the n -sum. Next, we apply the Poisson summation formula on the sum over n . This gives

$$\begin{aligned} &\sum_{n \in \mathbb{Z}} \chi_1\chi_2(n) e\left(\frac{\overline{M_3}mn}{qM_1}\right) \mathcal{J}(m, n, q) W\left(\frac{n}{N}\right) \\ &= \frac{N}{qM_1M_2} \sum_{n \in \mathbb{Z}} \mathfrak{C}(m, n, q) \mathcal{J}(m, n, q), \end{aligned}$$

where the character sum is given by

$$\mathfrak{C}(m, n, q) = \sum_{a \pmod{M_1M_2q}} \chi_1\chi_2(a) e\left(\frac{\overline{M_3}ma}{qM_1} + \frac{na}{qM_1M_2}\right)$$

and the integral is

$$\begin{aligned} \mathcal{J}(x, y, q) &= \int_{\mathbb{R}} \mathcal{J}(x, Nv, q) W(v) e\left(-\frac{Nvy}{qM_1M_2}\right) dv \\ &= \iint_{\mathbb{R}^2} h\left(\frac{q}{Q}, v-u\right) W(v) V(u) e\left(-\frac{Nux}{qM_1M_3} - \frac{Nvy}{qM_1M_2}\right) dudv. \end{aligned}$$

By repeated integration-by-parts we get that the tail $|n| \gg QM_1M_2M^\varepsilon/N$ makes a negligible contribution to the sum.

We impose the restriction that $Q \ll M_2$ with a sufficiently small implied constant. This is equivalent to having

$$(3.3) \quad N \ll M_1M_2^2$$

Under this condition $(q, M_2) = 1$, and therefore the character sum splits as

$$\sum_{a \pmod{qM_1}} \chi_1(a) e\left(\frac{\overline{M_3}ma}{qM_1}\right) e\left(\frac{\overline{M_2}na}{qM_1}\right) \sum_{b \pmod{M_2}} \chi_2(b) e\left(\frac{\overline{qM_1}nb}{M_2}\right),$$

which is

$$\varepsilon_2 \sqrt{\overline{M_2}} \chi_2(qM_1) \overline{\chi}_2(n) \sum_{a \pmod{qM_1}} \chi_1(a) e\left(\frac{(\overline{M_3}m + \overline{M_2}n)a}{qM_1}\right).$$

Suppose $q = q'M_1^r$ with $M_1 \nmid q'$. Then the remaining character sum splits into the product

$$\begin{aligned} &\sum_{a \pmod{q'}} e\left(\frac{(\overline{M_3}m + \overline{M_2}n)\overline{M_1}^{r+1}a}{q'}\right) \\ &\quad \times \sum_{b \pmod{M_1^{r+1}}} \chi_1(b) e\left(\frac{(\overline{M_3}m + \overline{M_2}n)\overline{q'}b}{M_1^{r+1}}\right). \end{aligned}$$

This product vanishes unless $q'M_1^r | M_2m + M_3n$ in which case we get

$$q'\varepsilon_1 \sqrt{\overline{M_1}} M_1^r \chi_1(q'M_2M_3) \overline{\chi}_1((M_2m + M_3n)/M_1^r).$$

We conclude that

$$\begin{aligned} &\sum_{n \in \mathbb{Z}} \chi_1 \chi_2(n) e\left(\frac{\overline{M_3}mn}{qM_1}\right) W\left(\frac{n}{N}\right) \mathcal{J}(m, n, q) = \frac{\varepsilon_1 \varepsilon_2 N}{\sqrt{M_1 M_2}} \\ &\sum_{\substack{|n| \ll QM_1M_2M^\varepsilon/N \\ q | M_2m + M_3n}} \chi_1(q'M_2M_3) \overline{\chi}_1((M_2m + M_3n)/M_1^r) \chi_2(qM_1) \overline{\chi}_2(n) \mathcal{J}(m, n, q) \\ &\quad + O_A(M^{-A}), \end{aligned}$$

where $r = v_{M_1}(q)$ is the M_1 -adic valuation of q and $q' = q/M_1^r$. Since $(m, q) = 1$, we also have $(n, q) = 1$. Consequently

$$S_\chi(N) = \frac{\eta c_Q N}{\sqrt{M}} \sum_{q \ll Q} \chi_1(q') \sum_{\substack{|n| \ll Q M_1 M_2 M^\varepsilon / N \\ |m| \ll Q M_1 M_3 M^\varepsilon / N \\ (m, q M_3) = (n, q M_2) = 1 \\ q | M_2 m + M_3 n}} \bar{\chi}_1((M_2 m + M_3 n)/M_1^r) \chi_2(q) \bar{\chi}_2(n) \chi_3(q) \bar{\chi}_3(m) \mathcal{J}(m, n, q) + O_A(M^{-A}),$$

with $|\eta| = 1$. At this stage, trivial estimation gives

$$S_\chi(N) \ll M^{1/2+\varepsilon},$$

which is just at the threshold. Any additional saving will yield a non-trivial bound for the character sum.

Observe that for $r \geq 1$ we are saving an extra M_1 by trivial estimation. Therefore, we just need to focus on the generic case $r = 0$. We consider

$$S_0 := \sum_{\substack{q \ll Q \\ (q, M_1) = 1}} \chi_1 \chi_2 \chi_3(q) \sum_{\substack{|n| \ll Q M_1 M_2 M^\varepsilon / N \\ |m| \ll Q M_1 M_3 M^\varepsilon / N \\ (m, q M_3) = (n, q M_2) = 1 \\ q | M_2 m + M_3 n}} \bar{\chi}_1(M_2 m + M_3 n) \bar{\chi}_2(n) \bar{\chi}_3(m) \mathcal{J}(m, n, q).$$

Then

$$(3.4) \quad S_\chi(N) = \frac{\eta c_Q N}{\sqrt{M}} S_0 + O\left(\frac{N M^\varepsilon}{M_1}\right).$$

3.3. Treatment of S_0 .

3.3.1. *Applying Cauchy's inequality.* From Cauchy's inequality we get

$$(3.5) \quad S_0 \ll M^\varepsilon \sqrt{\frac{Q M_1 M_2}{N}} \sqrt{T},$$

where T is given by

$$\sum_{|n| \ll Q M_1 M_2 M^\varepsilon / N} \left| \sum_{\substack{q \ll Q \\ (q, M_1 n) = 1}} \chi_1 \chi_2 \chi_3(q) \sum_{\substack{|m| \ll Q M_1 M_3 M^\varepsilon / N \\ (m, q M_3) = 1 \\ q | M_2 m + M_3 n}} \bar{\chi}_1(M_2 m + M_3 n) \bar{\chi}_3(m) \mathcal{J}(m, n, q) \right|^2.$$

Here we assumed that $M_3 < M_2$, otherwise we would have pulled out the m -sum rather than the n -sum. Any non-trivial bound for T will yield a non-trivial bound for the character sum. Introducing a dyadic partition of unity for the n -sum and opening the absolute square it suffices to consider the following sum

$$(3.6) \quad \sum_{\substack{q_1, q_2 \ll Q \\ (q_i, M_1) = 1}} \chi_1 \chi_2 \chi_3(q_1 \bar{q}_2) \sum_{\substack{|m_i| \ll Q M_1 M_3 M^\varepsilon / N \\ (m_i, q_i M_3) = 1}} \chi_3(\bar{m}_1 m_2) T(m_1, m_2, q_1, q_2)$$

where $T(m_1, m_2, q_1, q_2)$ is given by

$$\sum_{\substack{n \in \mathbb{Z} \\ n \equiv -M_2 \overline{M_3 m_i} \pmod{q_i}}} \overline{\chi}_1(M_2 m_1 + M_3 n) \chi_1(M_2 m_2 + M_3 n) \\ \times \mathcal{K}(m_1, m_2, n, q_1, q_2) U\left(\frac{n}{R}\right),$$

with

$$\mathcal{K}(x_1, x_2, y, q_1, q_2) = \mathcal{J}(x_1, y, q_1) \overline{\mathcal{J}(x_2, y, q_2)}.$$

Here U is a suitable smooth function with compact support and $R \ll Q M_1 M_2 M^\varepsilon / N$.

3.3.2. The third application of Poisson summation. We seek to get cancellation in $T(m_1, m_2, q_1, q_2)$. For this we at least need that the sum has enough points of summation or $Q M_1 M_2 / N \gg Q^2$, which is equivalent to

$$(3.7) \quad N \ll M_1 M_2^{2/3}.$$

We now apply Poisson summation to $T(m_1, m_2, q_1, q_2)$ with modulus $q_1 q_2 M_1$. This gives

$$T(m_1, m_2, q_1, q_2) = \frac{R}{q_1 q_2 M_1} \sum_{n \in \mathbb{Z}} \mathfrak{C}(m_1, m_2, n, q_1, q_2) \mathcal{L}(m_1, m_2, n, q_1, q_2)$$

where the character sum $\mathfrak{C}(m_1, m_2, n, q_1, q_2)$ is given by

$$\sum_{\substack{a \pmod{q_1 q_2 M_1} \\ a \equiv -M_2 \overline{M_3 m_i} \pmod{q_i}}} \overline{\chi}_1(M_2 m_1 + M_3 a) \chi_1(M_2 m_2 + M_3 a) e\left(\frac{na}{q_1 q_2 M_1}\right),$$

and the integral is given by

$$\mathcal{L}(x_1, x_2, z, q_1, q_2) = \int_{\mathbb{R}} \mathcal{K}(x_1, x_2, Ry, q_1, q_2) U(y) e\left(-\frac{Ryz}{q_1 q_2 M_1}\right) dy.$$

By repeated integration-by-parts we see that the integral $\mathcal{L}(m_1, m_2, n, q_1, q_2)$ is negligibly small if $|n| \gg Q^2 M_1 M^\varepsilon / R = N M^\varepsilon / R$. Hence

$$T(m_1, m_2, q_1, q_2) = \frac{R}{q_1 q_2 M_1} \sum_{|n| \ll N M^\varepsilon / R} \mathfrak{C}(m_1, m_2, n, q_1, q_2) \mathcal{L}(m_1, m_2, n, q_1, q_2) + O_A(M^{-A}).$$

Using the trivial bound $|\mathcal{L}(m_1, m_2, n, q_1, q_2)| \ll Q^2 / q_1 q_2$ which follows from (2.3), we conclude that

$$T(m_1, m_2, q_1, q_2) \ll \frac{R Q^2}{(q_1 q_2)^2 M_1} \sum_{|n| \ll N M^\varepsilon / R} |\mathfrak{C}(m_1, m_2, n, q_1, q_2)| + M^{-A}.$$

3.3.3. Bounds for $\mathfrak{C}(m_1, m_2, n, q_1, q_2)$ and $T(m_1, m_2, q_1, q_2)$. Since $M_1 \nmid q_1 q_2$, the character sum $\mathfrak{C}(m_1, m_2, n, q_1, q_2)$ splits as

$$\sum_{a \pmod{M_1}} \overline{\chi}_1(M_2 m_1 + M_3 a) \chi_1(M_2 m_2 + M_3 a) e\left(\frac{\overline{q_1 q_2} n a}{M_1}\right) \\ \times \sum_{\substack{b \pmod{q_1 q_2} \\ b \equiv -M_2 \overline{M_3 m_i} \pmod{q_i}}} e\left(\frac{\overline{M_1} n b}{q_1 q_2}\right).$$

The second sum has bound (q_1, q_2) , since there are at most (q_1, q_2) many terms due to the Chinese remainder theorem. On making the change of indices $x_1 = M_2 m_2 + M_3 a$, one sees that the first sum is equal to

$$\eta \mathfrak{S}_{\chi_1} (M_2(m_1 - m_2) \pmod{M_1}, \overline{q_1 q_2 M_3 n} \pmod{M_1})$$

for some η with $|\eta| = 1$. Here we recall that the character sum $\mathfrak{S}_{\chi}(m, n)$ is defined by (2.6) in Section 2.3.

At this point we need to apply bounds from Section 2.3. In view of Lemma 3, we have the following uniform bound

$$\mathfrak{C}(m_1, m_2, n, q_1, q_2) \ll (q_1, q_2) \sqrt{M_1(n, M_1)}.$$

However, for the zero frequency $n = 0$ we shall use the bound

$$|\mathfrak{C}(m_1, m_2, 0, q_1, q_2)| \leq (q_1, q_2)(m_1 - m_2, M_1).$$

We conclude that $T(m_1, m_2, q_1, q_2)$ is dominated by

$$\frac{(q_1, q_2) R Q^2}{(q_1 q_2)^2 M_1} \left((m_1 - m_2, M_1) + \sqrt{M_1} \sum_{0 \neq |n| \ll N M^\varepsilon / R} \sqrt{(n, M_1)} \right).$$

This gives

$$(3.8) \quad T(m_1, m_2, q_1, q_2) \ll \frac{(q_1, q_2) R Q^2 M^\varepsilon}{(q_1 q_2)^2 M_1} (m_1 - m_2, M_1) + \frac{(q_1, q_2) Q^2 N M^\varepsilon}{(q_1 q_2)^2 \sqrt{M_1}}.$$

3.3.4. Bound for S_0 . To bound T , we will now substitute the bound (3.8) in to (3.6) and estimate the remaining sums trivially. To estimate the contribution of the first term in (3.8), we observe that

$$\frac{R Q^2}{M_1} \sum_{q_1, q_2 \ll Q} \sum_{\substack{|m_i| \ll Q M_1 M_3 M^\varepsilon / N \\ (m_i, q_i M_3)=1}} \frac{(q_1, q_2)}{(q_1 q_2)^2} \ll \frac{\sqrt{M_1} M_2 M_3^2}{N^{3/2}} M^\varepsilon$$

and also

$$R Q^2 \sum_{q_1, q_2 \ll Q} \sum_{\substack{|m_i| \ll Q M_1 M_3 M^\varepsilon / N \\ (m_i, q_i M_3)=1 \\ M_1 | m_2 - m_1}} \frac{(q_1, q_2)}{(q_1 q_2)^2} \ll \frac{M_1 M_2 M_3}{N} \left(1 + \frac{M_3}{\sqrt{N M_1}} \right) M^\varepsilon.$$

To estimate the contribution of the second term in (3.8) towards T we evaluate

$$\frac{Q^2 N}{\sqrt{M_1}} \sum_{q_1, q_2 \ll Q} \sum_{\substack{|m_i| \ll Q M_1 M_3 M^\varepsilon / N \\ (m_i, q_i M_3)=1}} \frac{(q_1, q_2)}{(q_1 q_2)^2} \ll \sqrt{M_1} M_3^2 M^\varepsilon.$$

Inserting these bounds in (3.5) we obtain

$$(3.9) \quad S_0 \ll \left(\frac{\sqrt{M_1} M_2 M_3}{N} + \frac{M_1^{3/4} M_2 \sqrt{M_3}}{N^{3/4}} + \frac{\sqrt{M_1} M_2 M_3}{N^{1/4}} \right) M^\varepsilon.$$

3.4. Conclusion. Observe that the first term in (3.9) absorbs the error term in (3.4), and therefore we obtain from (3.4) the following bound for $S_\chi(N)$,

$$S_\chi(N) \ll_\varepsilon \left(M_2^{1/2} M_3^{1/2} + M_1^{1/4} M_2^{1/2} N^{1/4} + M_3^{1/2} N^{3/4} \right) M^\varepsilon$$

which matches with (1.2) in Theorem 1. Also note that our assumptions (3.1), (3.2), (3.3) and (3.7) produced the condition

$$M_1 \ll N \ll M_1 \min \{ M_2^{2/3}, M_3^2 \}.$$

Dividing the above bound for $S_\chi(N)$ by $\sqrt{N}$ and returning to (2.1), we see that $L(\frac{1}{2}, \chi)$ is bounded by

$$M^{1/4-\delta/2+\varepsilon} + \max_{M^{1/2-\delta} \ll N \ll M^{1/2}} \left(\left(\frac{M_2 M_3}{N} \right)^{1/2} + \left(\frac{M_1 M_2^2}{N} \right)^{1/4} + M_3^{1/2} N^{1/4} \right) M^\varepsilon$$

for any $\varepsilon > 0$ provided that $M_1 \ll N \ll M_1 \min \{ M_2^{2/3}, M_3^2 \}$ is satisfied for all $M^{1/2-\delta} \ll N \ll M^{1/2+\varepsilon}$. Therefore, in order to establish Corollary 1, one needs

$$(3.10) \quad \max_{M^{1/2-\delta} \ll N \ll M^{1/2}} \max \left\{ \frac{M_2 M_3}{N}, \frac{M_1^{1/2} M_2}{N^{1/2}}, M_3 N^{1/2}, M_1 \right\} \ll M^{1/2-\delta},$$

and $M^{1/2+\varepsilon} \ll M_1 \min \{ M_2^{2/3}, M_3^2 \}$. One can easily verify that the last bound is always satisfied when (3.10) is satisfied. Choosing $N = M^{1/2-\delta}$ for the first two terms in (3.10) and $N = M^{1/2}$ for the third term in (3.10), we obtain the conditions given by the inequalities in (1.3) in Corollary 1.

References

- [Bur62a] D. A. Burgess, *On character sums and L-series*, Proc. London Math. Soc. (3) **12** (1962), 193–206. MR 0132733 (24 #A2570)
- [Bur62b] ———, *On character sums and primitive roots*, Proc. London Math. Soc. (3) **12** (1962), 179–192. MR 0132732 (24 #A2569)
- [Bur63] ———, *On character sums and L-series. II*, Proc. London Math. Soc. (3) **13** (1963), 524–536. MR 0148626 (26 #6133)
- [DFI93] W. Duke, J. Friedlander, and H. Iwaniec, *Bounds for automorphic L-functions*, Invent. Math. **112** (1993), no. 1, 1–8. MR 1207474 (94c:11043)
- [Fu09] Lei Fu, *Weights of twisted exponential sums*, Math. Z. **262** (2009), no. 2, 449–472. MR 2504886 (2009m:14031)
- [FW09] Brooke Feigon and David Whitehouse, *Averages of central L-values of Hilbert modular forms with an application to subconvexity*, Duke Math. J. **149** (2009), no. 2, 347–410. MR 2541706 (2010m:11067)
- [HB96] D. R. Heath-Brown, *A new form of the circle method, and its application to quadratic forms*, J. Reine Angew. Math. **481** (1996), 149–206. MR 1421949 (97k:11139)
- [HM12] Roman Holowinsky and Ritabrata Munshi, *Level aspect subconvexity for Rankin-Selberg L-functions*, Automorphic Representations and L-Functions, Tata Inst. Fund. Res. Stud. Math., Tata Inst. Fund. Res., Mumbai, 2012, pp. 311–334.
- [HMQ14] Roman Holowinsky, Ritabrata Munshi, and Zhi Qi, *Hybrid subconvexity bounds for $L(\frac{1}{2}, \text{Sym}^2 f \otimes g)$* , arxiv:1401.6695 (2014).
- [HT14] Roman Holowinsky and Nicolas Templier, *First moment of Rankin-Selberg central L-values and subconvexity in the level aspect*, Ramanujan J. **33** (2014), no. 1, 131–155. MR 3142436
- [IK04] Henryk Iwaniec and Emmanuel Kowalski, *Analytic number theory*, American Mathematical Society Colloquium Publications, vol. 53, American Mathematical Society, Providence, RI, 2004. MR 2061214 (2005h:11005)

- [Mil14] Djordje Milićević, *Sub-Weyl subconvexity for Dirichlet L -functions to prime power moduli*, arXiv:1407.4100 **preprint** (2014).
- [MR12] Philippe Michel and Dinakar Ramakrishnan, *Consequences of the Gross-Zagier formulae: stability of average L -values, subconvexity, and non-vanishing mod p* , Number theory, analysis and geometry, Springer, New York, 2012, pp. 437–459.
- [Mun13a] Ritabrata Munshi, *The circle method and bounds for L -functions—III: t -aspect subconvexity for $GL(3)$ L -functions*, arXiv:1301.1007 **preprint** (2013).
- [Mun13b] ———, *The circle method and bounds for L -functions—IV: Subconvexity for twists of $GL(3)$ L -functions - B*, arXiv:1311.6120 **preprint** (2013).
- [Mun14a] ———, *The circle method and bounds for L -functions—I*, Math. Ann. **358** (2014), no. 1-2, 389–401. MR 3158002
- [Mun14b] ———, *The circle method and bounds for L -functions—II: Subconvexity for twists of $GL(3)$ L -functions*, American J. Math. **to appear** (2014).
- [Nel13] Paul D. Nelson, *Stable averages of central values of Rankin-Selberg L -functions: some new variants*, J. Number Theory **133** (2013), no. 8, 2588–2615. MR 3045204
- [RS94] Zeév Rudnick and Peter Sarnak, *The behaviour of eigenstates of arithmetic hyperbolic manifolds*, Comm. Math. Phys. **161** (1994), no. 1, 195–213. MR 1266075 (95m:11052)

DEPARTMENT OF MATHEMATICS, THE OHIO STATE UNIVERSITY, 100 MATH TOWER, 231 WEST 18TH AVENUE, COLUMBUS, OH 43210, USA

E-mail address: `holowinsky.1@osu.edu`

SCHOOL OF MATHEMATICS, TATA INSTITUTE OF FUNDAMENTAL RESEARCH, 1 HOMI BHABHA ROAD, COLABA, MUMBAI 400005, INDIA

E-mail address: `rmunshi@math.tifr.res.in`

DEPARTMENT OF MATHEMATICS, THE OHIO STATE UNIVERSITY, 100 MATH TOWER, 231 WEST 18TH AVENUE, COLUMBUS, OH 43210, USA

E-mail address: `qi.91@osu.edu`